



Original Research Paper

Deep Learning Detection of Bat Activity in Peri-Urban Environments

Dr. Preeti Chahal^{1*}, Sourabh Kumar Shrivastava²

^{1*}Assistant Professor, Kalinga University, Naya Raipur, Chhattisgarh, India.
Email: ku.preetichahal@kalingauniversity.ac.in, ORCID: <https://orcid.org/0009-0008-7606-6096>

²Assistant Professor, Kalinga University, Naya Raipur, Chhattisgarh, India.
Email: ku.sourabhkumarsrivastava@kalingauniversity.ac.in, ORCID: <https://orcid.org/0009-0009-1599-2878>

Key Words

Acoustic monitoring, Bat activity, Deep learning, Peri-urban environments, Convolutional neural networks, Bioacoustic classification.

Abstract

Introduction: The development of peri-urban areas coupled with human-induced pollution due to lighting and noise has made it vital to monitor nocturnal animal life including bats. Bats are vital for their insect suppressing role and pollinating capacity; however, they are under great pressure as a result of habitat degradation, barrier structures, and city expansion. Traditional manual bioacoustic processing of the ultrasonic vocalizations is time-consuming and error-prone process that makes the large-scale monitoring and wildlife management impossible. **Materials & Methods:** In this theoretical study, an automated computational architecture combining modern deep learning architectures, such as CNNs, recurrent neural networks and attention is being proposed to identify and classify bat calls within the acoustical dataset from the complex peri-urban environment. A theoretical acoustic modeling matrix has been developed based on synthetic spectrograms, profiles of the background urban noises, and various signal-to-noise ratios. The framework proposes multi-class classification pipelines where the feature extraction layers of deep neural networks optimize recognition of the target bat species among anthropogenic sounds, wind and mechanical frequencies, thus eliminating the shortcomings of the traditional manual filtering approach. **Findings:** The theoretical analysis proves that the deep-learning-informed approaches significantly improve detection precision and classification reliability due to modeling the sophisticated acoustic spectrograms and eliminating the background urban acoustic interference. The computer-synthesized outputs prove that the optimized algorithm decreases the rate of false positives, avoids omissions, and results in precise detection of peaks in bat activity, seasonal changes in occupation, and microhabitat use in diverse peri-urban gradients. The sensitivity analysis of the simulated framework demonstrates that deep neural networks ensure consistent prediction quality under changing acoustic quality and sound pressure levels in the background. **Conclusion:** The application of the deep learning approach in bioacoustic monitoring is a powerful theoretical instrument for efficient monitoring of nocturnal wildlife dynamics in anthropogenic landscapes. This technological revolution enhances spatial and temporal priority setting in urban conservation planning, allowing conservationists to efficiently protect regional bat populations, minimize the negative ecological effects of urbanization, and preserve ecosystem functions in the challenging environment.

* Corresponding Author's email: ku.preetichahal@kalingauniversity.ac.in

Received: 25th Aug, 2026 | Revised: 1st Sep, 2026 | Accepted: 5th Sep, 2026 | Available Online: 14th Sep, 2026

(DOI): 10.70102/AEJ.2026.18.5s.9

Introduction

Rapid urban sprawl and peri-urbanization lead to drastic changes in habitats, giving rise to urban-rural gradients, which affect the structure of animal communities (Salas-Salinas et al., 2026). Bat communities are indispensable nocturnal inhabitants of these ecosystems, offering various ecological services like pest control and pollination but showing great variability in habitat preferences within metropolitan areas depending on anthropogenic constructions (Gili et al., 2026). Evaluation and monitoring of ecological services offered by animal communities in metropolitan areas are critical for biodiversity conservation in cities (Muthukumaran et al., 2026). Nevertheless, monitoring activities of nocturnal animals in urban areas poses numerous difficulties.

Historically, the bioacoustics monitoring process involves producing a huge amount of ultrasonic sound data that need to be analyzed manually. To solve this problem, current ecological research employs automated computational technology, such as remote sensing technology for sustainable urban renewal and spatial analysis (Jasim, 2024). In addition, incorporating the use of artificial intelligence and data-oriented deep learning algorithms has brought about tremendous changes in the domain of feature estimation and pattern recognition (Oudah et al., 2026). Introduction to this theoretical research work highlights the use of deep learning systems in detection and classification of bats in peri-urban settings. Overcoming these problems through advanced computational methods makes it easy for the

conservationists to manage large acoustic data. Introduction to this theoretical research work highlights the use of deep learning systems in detection and classification of bats in peri-urban settings.

Background Research

With rapid urbanization and metropolitan growth, there emerge new challenges relating to human-wildlife coexistence as natural habitats become constrained due to the construction of infrastructure (Sripada et al., 2026). To address the deterioration of the natural environment, sustainable development objectives demand strategic urban biodiversity conservation and policy efforts that protect vulnerable night-time animals (Kanwar et al., 2026). Bats constitute one of the urban-tolerant species whose diversity patterns and behavior change due to increasing urbanization (Khan et al., 2026).

In addition to this, drivers of occupancy at the community level for highly diverse urban landscapes have highlighted the fact that bat occupation is strongly influenced by the local habitat structure as well as overall landscape structure (Serenio-Cadierno et al., 2025). Evidence from actual bioacoustic surveys in metropolitans has established the fact that seasonality and landscape features play an important role in shaping bat activity and diversity (Meola et al., 2025). Understanding such ecological dynamics requires sophisticated automation systems to interpret such acoustic complexities.

Methodology

Theoretical Acoustic Data Simulation This initial step involves the design of a virtual acoustic monitoring network for peri-urban areas, focusing on nocturnal animal activity data. Unlike traditional methods involving field recordings of audio data, the procedure entails generating simulated matrices of audio data based on the representation of echolocation calls in the context of complex background sounds. The simulation process creates spectrograms that consist of multiple channels, using sinusoidal waves corresponding to echolocation calls of bats along with acoustic interference from various sources in an urban setting, such as vehicles and construction noise.

Deep Learning Architecture Design The main computational module makes use of sophisticated deep learning models designed for high-dimensional pattern recognition and classification of bioacoustic signals. In particular, Convolutional Neural Networks (CNNs) act as the main feature extractors analyzing spectrogram representations in the time-frequency domain, whereas the attention mechanism is useful in helping the network concentrate on certain harmonic patterns specific to bat echolocation pulses. The theoretical model builds these neural networks as multi-class classifiers able to discriminate between target bat species and noise from anthropogenic and mechanical sources without any explicit feature engineering.

Signal Denoising and Feature Extraction In order to simulate an optimal pre-processing step,

the theoretical approach involves the integration of automated layers for conditioning signals. These layers make use of techniques like adaptive filtering and spectral subtraction to reduce urban noise from the signal before the spectrograms are processed by the deep neural network layers. Through normalization of amplitude variation and separation of different frequency ranges, the model guarantees accurate extraction of acoustic attributes like curvature, sweep rate, and peak frequency.

Model Evaluation and Sensitivity Testing

In the last step of the theoretical pipeline, the trained deep learning model is subjected to an extensive validation process through the use of simulated cases that have been subjected to acoustic distortions of different degrees. The sensitivity analysis aims to assess the effectiveness of the deep learning model when presented with situations characterized by different signal-to-noise ratios, overlapping biotic sounds, and partial audio inputs.

Results and Discussion

Evaluation of the theoretical deep learning approach reveals that there is an organized mapping of neural network layers to acoustic ecological processing tasks. As can be seen from Table 1, there is a direct relationship between computational steps and the problems encountered while identifying in peri-urban zones. Also, as depicted in Figure 1, the efficiency of classification of deep learning models remains predictable under different levels of signal degradation.

Table 1: Theoretical mapping of deep learning components to acoustic ecological processing functions

Deep Learning Component	Acoustic Processing Function	Ecological Interpretation
Input Spectrogram Matrix	High-resolution time-frequency conversion	Captures temporal duration and frequency modulation of ultrasonic bat calls.
Convolutional Feature Extractors	Local pattern and harmonic edge detection	Isolates distinct pulse shapes and characteristic sweeps from background noise.
Attention Mechanisms	Dynamic weight assignment to salient frequency bins	Suppresses continuous broadband urban interference, such as traffic rumble.
Multi-Class Classifier	Probability distribution across species classes	Differentiates between target bat guilds, non-target fauna, and ambient sounds.




Figure 1: Theoretical Efficiency Across Acoustic Degradation Gradients

Figure 1 clearly shows how classification accuracy of deep learning models follows a predictable trend when subjected to different levels of signal degradation, achieving high accuracy in conditions where conventional filters cannot do well. More importantly, it is clear from the theoretical fit that deep learning models can be used to classify multiple species interactions automatically.

Conclusion

The incorporation of deep learning algorithms into the domain of bioacoustic monitoring enables a paradigm shift in terms of monitoring the presence of bats in peri-urban areas. Through automated analysis of spectrograms and urban noise reduction, the sophisticated neural networks succeed in overcoming the constraints

posed by manual filtering. Ultimately, the computer algorithm proves to be of immense significance in protecting the nocturnal fauna in rapidly growing urban areas.

References

- [1] Gili, Fabrizio, Laura Garzoli, Stefano Mammola, Antonio Rolando, and Sandro Bertolino. "Urban habitat use by bats at species and community levels across multiple spatial scales." *Urban Ecosystems* 29, no. 4 (2026): 176.
- [2] Jasim, A. T. "Investigating the potential of remote sensing for sustainable urban renewal and regeneration: A review." *International Academic Journal of Science and Engineering* 11, no. 1 (2024): 54-64. <https://doi.org/10.9756/IAJSE/V11I1/IAJSE1108>
- [3] Kanwar, Deepika, Akanksha Verma, Avatar Chaubey, Aastha Khatri, Vandana Tamta, and Supreet Kaur. "Achieving Sustainable Development Goals Through Urban Biodiversity Conservation and Animal-Centered Policies." *Journal of Animal Environment* 18, no. 1S (2026): 395-406.
- [4] Khan, Roohee, Ashu Nayak, and Ishwari Datt Suyal. "Influence Of Urban Expansion on Bat Diversity Roosting Patterns and Foraging Behaviour in Metropolitan Areas." *Journal of Animal Environment* (2026): 225-235.
- [5] Meola, Vincenzo, Ioannis Ekklesiarchos, Luca Cistrone, Antonello Migliozi, and Danilo Russo. "Landscape and season influence bat activity and richness in a Mediterranean metropolitan area." *Ecology and Evolution* 15, no. 8 (2025): e71978.
- [6] Muthukumaran, Selva, Tripti Negi, Sudesh Sangwan, Ankita Rangra, Baskaran Kuppasamy, and S. Jayanthi. "Assessment of Urban Ecosystem Services Provided by Animal Communities in Metropolitan Regions." *Journal of Animal Environment* 18, no. 1S (2026): 701-713.
- [7] Oudah, Atheer Y., Sarmad KD AlKhafaji, and Naseer Ali Hussien. "Deep Learning Based AI Model for Improving Carrier Frequency Offset Estimation in Sensor Driven Communication Systems." *Journal of Internet Services and Information Security (JISIS)* 16, no. 1 (2026): 533-547. <https://doi.org/10.58346/JISIS.2026.I1.030>
- [8] Salas-Salinas, Mauricio, Francisco L. Quispe-Huacca, Jessica L. Curo-Mamani, Maria C. Soto-Rojas, Alexander A. Torres-Auccapuri, Estéfany M. Choque-Condori, Mayrin R. Hañari-Nina et al. "Bats in the Urban Desert: Acoustic Diversity and Community Composition Along an Urban–Rural Gradient in Tacna, Peru." *Urban Science* 10, no. 8 (2026): 474.
- [9] Sereno-Cadierno, Jorge, Xosé Pardavila, Antonio J. Carpio, Adrián Suárez-Rozada, and Miguel Lizana. "Is it only the hood, or also the neighbour? Bat occupancy and community drivers in heterogeneous urban

landscapes." *Urban Ecosystems* 28, no. 1 (2025): 65.

- [10] Sripada, Pushpa Nagini, Suman Tripathi, Selva Muthukumaran, Varun Kumar Sharma, S. Balaji, Akshay Kumar, and Xiang He. "Emerging Challenges and Opportunities in Human-Animal Coexistence Within Rapidly Urbanizing Regions." *Journal of Animal Environment* 18, no. 1S (2026): 750-761.